

HEDIS® 2014 MQIC MEASURES SUMMARY LISTING FOR ANNUAL PERFORMANCE REPORTING

DIABETES

HEDIS® Measures:

1. Comprehensive Diabetes Care (CDC):

Percentage of members 18-75 years of age with diabetes (type 1 and type 2) who had each of the following:

- a. HbA1c Testing
- b. HbA1c Poor Control (>9.0%)
- c. HbA1c Control (<8.0%)
- d. HbA1c Control (<7.0%) for a selected population (Additional exclusion criteria are required for this indicator)
- e. Eye Exam (retinal) performed
- f. LDL-C Screening
- g. LDL-C Control (<100 mg/dL)
- h. Medical Attention for Nephropathy
- i. Blood Pressure Control (<140/80 mm Hg)
- j. Blood Pressure Control (<140/90 mm Hg)

Non-HEDIS Measures:

1. Statin Therapy for Persons with Diabetes

Members ages 40-75 with at least one statin prescription filled during the measurement year.

ASTHMA

HEDIS® Measures:

1. Use of Appropriate Medications for People with Asthma (ASM):

Percent members 5-64 years of age who were identified as having persistent asthma and who were appropriately prescribed medication during the measurement year.

- a. 5-11 years
- b. 12-18 years
- c. 19-50 years
- d. 51-64 years
- e. Total (sum of the four numerators divided by sum of the four denominators)

Non-HEDIS Measures:

1. Periodic Assessment (5-64 Years Old)

Members who had at least one preventive/ambulatory health services visit within the measurement year with a PCP, pulmonologist, or allergist.

2. Emergency Department Visits per 1000 Members with Asthma and Being Treated for a Primary Diagnosis of Asthma (5-64 Years Old)

HYPERTENSION

HEDIS® Measure:

1. Controlling High Blood Pressure (CBP):

Percent members 18-85 years of age who had a diagnosis of HTN and whose BP was adequately controlled (<140/90 mm Hg) during the measurement year.

Non-HEDIS Measure:

1. Percent Members with any Diagnosis of Hypertension

ACUTE PHARYNGITIS

HEDIS® Measure:

1. Appropriate Testing for Children With Pharyngitis (CWP):

Percent children 2-18 years of age diagnosed with pharyngitis, dispensed an antibiotic and received a group A streptococcus test for the episode.

ROUTINE PRENATAL AND POSTNATAL CARE

HEDIS® Measures:

1. Prenatal and Postpartum Care (PPC):

The percentage of deliveries of live births between November 6 of the year prior to the measurement year and November 5 of the measurement year. For these women, the measure assesses the following facets of prenatal and postpartum care:

- a. Timeliness of Prenatal Care: percent deliveries that received a prenatal care visit as a member of the organization in the first trimester *or* within 42 days of enrollment in the organization.
 - b. Postpartum Care: percent deliveries that had a postpartum visit on or between 21 and 56 days after delivery.
-

CHRONIC KIDNEY DISEASE

Non-HEDIS Measure:

1. Percent members in HEDIS® Diabetes or Hypertension populations who had a serum creatinine test (necessary for GFR)

MANAGEMENT OF ACUTE LOW BACK PAIN

HEDIS® Measure:

1. Use of Imaging Studies for Low Back Pain (LBP):

Percent members with a primary diagnosis of low back pain who did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of diagnosis.

PREVENTIVE SERVICES FOR INFANTS AND CHILDREN (BIRTH – 24 MONTHS)

HEDIS® Measures:

1. Childhood Immunization Status (CIS):

Percent of children two years of age who had four DTaP, three IPV, one MMR, three Hib, three Hepatitis B, one VZV, four Pneumococcal conjugate, one Hepatitis A, two or three Rotavirus (number varies based on which vaccine is given), and two Influenza vaccines by their second birthday. The measure calculates a rate for each vaccine and nine separate combination rates, see Table 1.

Table 1 – Combination Vaccinations

Combination	DTaP	IPV	MMR	Hib	Hep B	VZV	PCV	Hep A	RV	Influenza
Combination 2*	X	X	X	X	X	X				
Combination 3*	X	X	X	X	X	X	X			
Combination 4	X	X	X	X	X	X	X	X		
Combination 5	X	X	X	X	X	X	X		X	
Combination 6	X	X	X	X	X	X	X			X
Combination 7	X	X	X	X	X	X	X	X	X	
Combination 8	X	X	X	X	X	X	X	X		X
Combination 9	X	X	X	X	X	X	X		X	X
Combination 10*	X	X	X	X	X	X	X	X	X	X

*MQIC health plans report all CIS measures, including Combos 2-10, but only Combos 2, 3 and 10 results are reported on the MQIC Annual Performance Measurement Report.

2. Well-Child Visits in the First 15 Months of Life (W15):

Percent members who turned 15 months old during the measurement year and who had **six** or more well-child visits with a PCP during their first 15 months of life.

3. Lead Screening in Children (LSC) [Medicaid]

Percent of children 2 years of age who had one or more capillary or venous lead blood test for lead poisoning by their second birthday.

PREVENTIVE SERVICES FOR CHILDREN AND ADOLESCENTS (AGES 2 - 21)

HEDIS® Measures:

1. Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life (W34):

Percent members 3-6 years of age who had one or more well-child visits with a PCP during the measurement year.

2. Adolescent Well-Care Visits (AWC):

Percent enrolled members 12-21 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement year.

3. Chlamydia Screening in Women (16 – 20 years) (CHL):

Percentage of women 16-20 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.

4. Immunizations for Adolescents (IMA):

Percent of adolescents 13 years of age who had one dose of meningococcal vaccine and one tetanus, diphtheria toxoids and acellular pertussis vaccine (Tdap) or one tetanus, diphtheria toxoids vaccine (Td) by their 13th birthday. This measure calculates a rate for each vaccine and one combination rate.

5. Human Papillomavirus Vaccine for Female Adolescents (HPV):

Percent of female adolescents 13 years of age who had three doses of the human papillomavirus (HPV) vaccine by their 13th birthday.